

**Lattice Degeneration** is a condition that involves abnormal thinning of the peripheral retina, which is the tissue that lines the back wall of the eye and is critical for maintaining good vision. When lattice degeneration is present, the retina is more vulnerable to developing tears, breaks, or holes that could ultimately lead to a visually debilitating condition called a **retinal detachment**. For this reason, once diagnosed lattice degeneration should be closely monitored.

Clinically, lattice degeneration is characterized by oval or straight patches of thinned retina, sometimes accompanied by pigment clumps or a crosshatching pattern formed by **sclerotic vessels** (Figure 1). Lattice may be found in only one eye, but often is present in both. There may be just one lesion or clusters of many. Lattice is often located along the outer border of the retina.


**Causes and Risk Factors:** Lattice degeneration occurs in 8% to 10% of the general population and its cause is not fully understood. There does not appear to be a correlation in incidence by gender or race. And while this condition does not follow a definitive inheritance pattern, it frequently clusters within families. It is most commonly found in patients with myopia (nearsightedness), but lattice-like lesions are also seen in rare diseases such as **Stickler syndrome**, **Ehlers-Danlos**, and **Marfan syndrome**. There is no prevention or cure for lattice degeneration.

**Symptoms and Diagnostic Testing:** Lattice degeneration itself does not cause symptoms, so the only way to diagnose the condition is with a dilated fundus examination by an eye care provider. A dilated **fundus** examination is done by administering dilating eye drops in your eyes to expand the pupil so that the retina can be carefully evaluated. Dilating drops will cause your vision to be blurry for several hours before returning to normal.


The eye care specialist will then use a headlight and a special lens to perform the exam. Depending on the nature of your findings, the doctor may also examine your retina while exerting a slight amount of pressure around your eye (scleral depression). In general, no imaging tests are necessary to diagnose this disease, although some providers may obtain wide-angle photographs of your retinas to assist in monitoring your condition.

*continued next page*

#### WHAT IS THE RETINA?



**THE RETINA** is a thin layer of light-sensitive nerve tissue that lines the back of the eye (or vitreous) cavity. When light enters the eye, it passes through the iris to the retina where images are focused and converted to electrical impulses that are carried by the optic nerve to the brain resulting in sight.



**Figure 1.**

Photograph of the peripheral retina demonstrates an area of lattice degeneration. Note the retinal thinning, which is characterized by a color change. There are also pigment clumps and crosshatching of sclerotic vessels, two features commonly observed in lattice degeneration.

Hamid Ahmadi, MD, Labbafinejad Medical Center. Retina Image Bank 2015; Image 25882. © 2018, American Society of Retina Specialists.

## Lattice Degeneration *continued from previous page*

**Treatment and Prognosis:** Lattice is thought to be minimally progressive and fortunately, the vast majority of patients with this condition never develop a retinal tear or detachment. Therefore, in general, no treatment is needed for lattice degeneration aside from regular monitoring with dilated fundus examinations.

In rare circumstances, some physicians may perform preventive laser therapy or **cryotherapy** (a type of freezing treatment) to strengthen the peripheral retina in the areas where it is weak. Your doctor will discuss the risks and benefits of doing so with you if he or she thinks it is worth considering. It is not known if these interventions are effective in preventing retinal complications.

Rarely, lattice degeneration can lead to complications such as a retinal tear or detachment. Symptoms to look out for include blurry vision, a curtain obscuring part of the outer visual field, flashing lights, and new floaters. If any of these occur, you should seek prompt attention from an ophthalmologist or retina specialist.

In those who do develop a retinal tear or detachment, treatment will be performed, typically by a retina specialist. This treatment can range from a laser treatment completed in the office to surgery in the operating room, depending on the severity of the condition.

**Conclusion:** In summary, lattice degeneration is a relatively common condition that affects the retina, especially in nearsighted people. While its presence increases the risk of a retinal tear or detachment, the vast majority of patients will never experience symptoms or complications from their lattice degeneration.

If you are diagnosed with lattice degeneration, the most important actions you can take to protect the health of your eyes are to visit your eye care provider on a regular basis and know the symptoms of a retinal tear or detachment so you can seek prompt treatment if needed. ●

### WHAT TO LOOK OUT FOR

**Lattice degeneration itself does not cause symptoms. However, in rare cases it can lead to a retinal tear or detachment, which are very serious conditions that require prompt attention from an ophthalmologist or retina specialist.**

**Symptoms include:**

- Blurred vision
- Flashing lights
- Floaters
- Curtain obscuring part of your peripheral visual field

If you experience any of these, seek prompt ophthalmic care. ●

### THANK YOU TO THE RETINA HEALTH SERIES AUTHORS

- Sophie J. Bakri, MD
- Audina Berrocal, MD
- Antonio Capone, Jr., MD
- Netan Choudhry, MD, FRCS-C
- Thomas Ciulla, MD, MBA
- Pravin U. Dugel, MD
- Geoffrey G. Emerson, MD, PhD
- Roger A. Goldberg, MD, MBA
- Darin R. Goldman, MD
- Dilraj Grewal, MD
- Larry Halperin, MD
- Vi S. Hau, MD, PhD
- Suber S. Huang, MD, MBA
- G. Baker Hubbard, MD
- Mark S. Humayun, MD, PhD
- Peter K. Kaiser, MD
- M. Ali Khan, MD
- Anat Loewenstein, MD
- Mathew J. MacCumber, MD, PhD
- Maya Maloney, MD
- Hossein Nazari, MD
- Oded Ohana, MD, MBA
- George Parlitsis, MD
- Jonathan L. Prenner, MD
- Gilad Rabina, MD
- Carl D. Regillo, MD, FACS
- Naryan Sabherwal, MD
- Sherveen Salek, MD
- Andrew P. Schachat, MD
- Michael Seider, MD
- Eduardo Uchiyama, MD
- Allen Z. Verne, MD
- Christina Y. Weng, MD, MBA
- Yoshihiro Yonekawa, MD

### EDITOR

John T. Thompson, MD

### MEDICAL ILLUSTRATOR

Tim Hengst

### Clinical Terms *(appearing green within fact sheet text)*

**Cryotherapy:** A technique in which a pencil-like probe is placed on the white of the eye (sclera). The tip of the probe becomes very cold and can freeze structures inside the eye. This can be used to create a seal around a retinal tear.

**Ehlers-Danlos:** A genetic connective tissue disease where joints are hyper-extendable and skin is abnormally elastic. This disease is associated with an increased risk of lattice degeneration and retinal detachment.

**Fundus:** The back of the eye where the retina, macula, vitreous, choroid, and optic nerve are located.

**Marfan syndrome:** A connective tissue disease where those affected are abnormally tall with long limbs, fingers and toes. Some with this disease develop life-threatening problems with their aorta and have an increased risk of lattice degeneration and retinal detachment.

**Retinal detachment:** A condition where the retina separates from the back of the eye cavity. This may be caused by vitreous gel or fluid leaking through a retinal tear or hole and collecting under the retina, causing it to separate from the tissue around it.

**Sclerotic vessels:** Blood vessels that are often thin and white rather than red due to lack of blood flow or tissue damage. These types of vessels are often found in patches of lattice degeneration.

**Stickler syndrome:** A genetic disorder characterized by flattened nose and cheekbones, problems with the skeletal system, hearing loss and lattice degeneration. The vitreous jelly is abnormally liquified which can lead to early onset retinal detachment.